

# Integration by Parts

## Question Paper 1

|                   |                      |
|-------------------|----------------------|
| <b>Level</b>      | A Level              |
| <b>Subject</b>    | Maths                |
| <b>Exam Board</b> | OCR - MEI            |
| <b>Module</b>     | Core 3               |
| <b>Topic</b>      | Calculus             |
| <b>Sub Topic</b>  | Integration by Parts |
| <b>Booklet</b>    | Question Paper 1     |

**Time Allowed:** 62 minutes

**Score:** /51

**Percentage:** /100

**Grade Boundaries:**

| A*   | A     | B   | C     | D     | E   | U    |
|------|-------|-----|-------|-------|-----|------|
| >85% | 77.5% | 70% | 62.5% | 57.5% | 45% | <45% |

1 Find the exact value of  $\int x^3 \ln x \, dx$ . [5]

2 Fig. 8 shows the curve  $y = f(x)$ , where  $f(x) = \frac{x}{\sqrt{2+x^2}}$




Fig. 8

(i) Show algebraically that  $f(x)$  is an odd function. Interpret this result geometrically. [3]

(ii) Show that  $f'(x) = \frac{2}{(2+x^2)^{\frac{3}{2}}}$ . Hence find the exact gradient of the curve at the origin. [5]

(iii) Find the exact area of the region bounded by the curve, the  $x$ -axis and the line  $x = 1$ . [4]

(iv) (A) Show that if  $y = \frac{x}{\sqrt{2+x^2}}$ , then  $\frac{1}{y^2} = \frac{2}{x^2} + 1$ . [2]

(B) Differentiate  $\frac{1}{y^2} = \frac{2}{x^2} + 1$  implicitly to show that  $\frac{dy}{dx} = \frac{2y^3}{x^3}$ . Explain why this expression cannot be used to find the gradient of the curve at the origin. [4]

3 Evaluate  $\int_0^3 x(x+1)^{-\frac{1}{2}} \, dx$ , giving your answer as an exact fraction. [5]

4 Show that  $\int_0^{\frac{\pi}{2}} x \cos^{\frac{1}{2}} x \, dx = \frac{\sqrt{2}}{2} \pi + 2\sqrt{2} - 4$ . [5]

5 Fig. 8 shows the curve  $y = \frac{x}{\sqrt{x-2}}$ , together with the lines  $y = x$  and  $x = 11$ . The curve meets these lines at P and Q respectively. R is the point (11, 11).




Fig. 8

(i) Verify that the  $x$ -coordinate of P is 3. [2]

(ii) Show that, for the curve,  $\frac{dy}{dx} = \frac{x-4}{2(x-2)^{\frac{3}{2}}}$ .

Hence find the gradient of the curve at P. Use the result to show that the curve is **not** symmetrical about  $y = x$ . [7]

(iii) Using the substitution  $u = x - 2$ , show that  $\int_3^{11} \frac{x}{\sqrt{x-2}} \, dx = 25\frac{1}{3}$ .

Hence find the area of the region PQR bounded by the curve and the lines  $y = x$  and  $x = 11$ . [9]