


**Mathematics (MEI)**

Advanced GCE 4753

Methods for Advanced Mathematics (C3)

**Mark Scheme for June 2010**

---

|   |   |  |   |
|---|---|--|---|
| <p><b>1</b></p> $\int_0^{\pi/6} \cos 3x \, dx = \left[ \frac{1}{3} \sin 3x \right]_0^{\pi/6}$ $= \frac{1}{3} \sin \frac{\pi}{2} - 0$ $= 1/3$  | <p>M1<br/>B1<br/>A1cao<br/>[3]</p>            | <p><math>k \sin 3x, k &gt; 0, k \neq 3</math><br/><math>k = (\pm)1/3</math><br/>0.33 or better</p>   | <p>or M1 for <math>u = 3x \Rightarrow \int \frac{1}{3} \cos u \, du</math> condone <math>90^\circ</math> in limit<br/>or M1 for <math>\left[ \frac{1}{3} \sin u \right]</math><br/>so: <math>\sin 3x</math>: M1B0, <math>-\sin 3x</math>: M0B0,<br/><math>\pm 3 \sin 3x</math>: M0B0, <math>-1/3 \sin 3x</math>: M0B1</p> |
| <p><b>2</b></p> <p><math>fg(x) =  x+1 </math>      <math>gf(x) =  x +1</math></p>    | <p>B1 B1<br/><br/>B1<br/>B1<br/>[4]</p>       | <p>soi from correctly-shaped graphs (i.e. without intercepts)<br/><br/>graph of <math> x+1 </math> only<br/>graph of <math> x +1</math></p>    | <p>but must indicate which is which<br/>bod gf if negative <math>x</math> values are missing<br/><br/>'V' shape with <math>(-1, 0)</math> and <math>(0, 1)</math> labelled<br/>'V' shape with <math>(0, 1)</math> labelled <math>(0, 1)</math></p>  |
| <p><b>3(i)</b></p> $y = (1+3x^2)^{1/2}$ $\Rightarrow \frac{dy}{dx} = \frac{1}{2}(1+3x^2)^{-1/2} \cdot 6x$ $= 3x(1+3x^2)^{-1/2}$   | <p>M1<br/>B1<br/>A1<br/>[3]</p>               | <p>chain rule<br/><math>\frac{1}{2} u^{-1/2}</math><br/>o.e., but must be '3'</p>  | <p>can isw here</p>   |
| <p><b>(ii)</b></p> $y = x(1+3x^2)^{1/2}$ $\Rightarrow \frac{dy}{dx} = x \cdot \frac{3x}{\sqrt{1+3x^2}} + 1 \cdot (1+3x^2)^{1/2}$ $= \frac{3x^2 + 1 + 3x^2}{\sqrt{1+3x^2}}$ $= \frac{1+6x^2}{\sqrt{1+3x^2}} *$ | <p>M1<br/>A1ft<br/><br/>M1<br/>E1<br/>[4]</p> | <p>product rule<br/>ft their <math>dy/dx</math> from (i)<br/><br/>common denominator or factoring<br/><math>(1+3x^2)^{-1/2}</math><br/>www</p> | <p>must show this step for M1 E1</p>  |

|  |  |  |  |
|--|--|--|--|
| <p><b>4</b> <math>p = 100/x = 100x^{-1}</math><br/> <math>\Rightarrow dp/dx = -100x^{-2} = -100/x^2</math></p> <p><math>dp/dt = dp/dx \times dx/dt</math><br/> <math>dx/dt = 10</math></p> <p>When <math>x = 50</math>, <math>dp/dx = (-100/50^2)</math><br/> <math>\Rightarrow dp/dt = 10 \times -0.04 = -0.4</math></p>  | <p>M1<br/>A1<br/>M1<br/>B1<br/>M1dep<br/>A1cao<br/>[6]</p>                 | <p>attempt to differentiate<br/> <math>-100x^{-2}</math> o.e.<br/> o.e. soi<br/> soi<br/> substituting <math>x = 50</math> into their <math>dp/dx</math> dep 2<sup>nd</sup> M1<br/> o.e. e.g. decreasing at 0.4</p>  | <p>condone poor notation if chain rule correct<br/> or <math>x = 50 + 10t</math> B1<br/> <math>\Rightarrow P = 100/x = 100/(50 + 10t)</math><br/> <math>\Rightarrow dP/dt = -100(50 + 10t)^{-2} \times 10 = -1000/(50 + 10t)^2</math> M1<br/> A1<br/> When <math>t = 0</math>, <math>dP/dt = -1000/50^2 = -0.4</math> A1</p>         |
| <p><b>5</b> <math>y^3 = xy - x^2</math><br/> <math>\Rightarrow 3y^2 dy/dx = x dy/dx + y - 2x</math></p> <p><math>\Rightarrow 3y^2 dy/dx - x dy/dx = y - 2x</math><br/> <math>\Rightarrow (3y^2 - x) dy/dx = y - 2x</math><br/> <math>\Rightarrow dy/dx = (y - 2x)/(3y^2 - x) *</math></p> <p>TP when <math>dy/dx = 0 \Rightarrow y - 2x = 0</math><br/> <math>\Rightarrow y = 2x</math><br/> <math>\Rightarrow (2x)^3 = x \cdot 2x - x^2</math><br/> <math>\Rightarrow 8x^3 = x^2</math><br/> <math>\Rightarrow x = 1/8</math> *(or 0)</p> | <p>B1<br/>B1<br/><br/>M1<br/>E1<br/><br/>M1<br/>M1<br/><br/>E1<br/>[7]</p> | <p><math>3y^2 dy/dx</math><br/> <math>x dy/dx + y - 2x</math></p> <p>collecting terms in <math>dy/dx</math> only</p> <p>or <math>x = 1/8</math> and <math>dy/dx = 0 \Rightarrow y = 1/4</math><br/> or <math>(1/4)^3 = (1/8)(1/4) - (1/8)^2</math><br/> or verifying e.g. <math>1/64 = 1/64</math></p> | <p>must show '<math>x dy/dx + y</math>' on one side</p> <p>or <math>x = 1/8 \Rightarrow y^3 = (1/8)y - 1/64</math> M1<br/> verifying that <math>y = 1/4</math> is a solution (must show evidence*) M1<br/> <math>\Rightarrow dy/dx = (1/4 - 2(1/8))/(...) = 0</math> E1<br/> *just stating that <math>y = 1/4</math> is M1 M0 E0</p> |
| <p><b>6</b> <math>f(x) = 1 + 2 \sin 3x = y \quad x \leftrightarrow y</math><br/> <math>x = 1 + 2 \sin 3y</math><br/> <math>\Rightarrow \sin 3y = (x - 1)/2</math><br/> <math>\Rightarrow 3y = \arcsin [(x - 1)/2]</math><br/> <math>\Rightarrow y = \frac{1}{3} \arcsin \left[ \frac{x-1}{2} \right]</math> so <math>f^{-1}(x) = \frac{1}{3} \arcsin \left[ \frac{x-1}{2} \right]</math></p> <p>Range of <math>f</math> is <math>-1</math> to <math>3</math><br/> <math>\Rightarrow -1 \leq x \leq 3</math></p>                            | <p>M1<br/>A1<br/>A1<br/><br/>A1<br/><br/>M1<br/>A1<br/>[6]</p>             | <p>attempt to invert</p> <p>must be <math>y = \dots</math> or <math>f^{-1}(x) = \dots</math></p> <p>or <math>-1 \leq (x - 1)/2 \leq 1</math><br/> must be '<math>x</math>', not <math>y</math> or <math>f(x)</math></p>  | <p>at least one step attempted, or reasonable attempt at flow chart inversion</p> <p>(or any other variable provided same used on each side)</p> <p>condone '&lt;'s for M1<br/> allow unsupported correct answers; <math>-1</math> to <math>3</math> is M1 A0</p>  |
| <p><b>7</b> (A) True , (B) True , (C) False<br/> Counterexample, e.g. <math>\sqrt{2} + (-\sqrt{2}) = 0</math></p>  | <p>B2,1,0<br/>B1<br/>[3]</p>   |  |  |

|  |   |   |  |
|--|---|---|--|
| <b>8(i)</b> When $x = 1$ , $y = 3 \ln 1 + 1 - 1^2 = 0$   | E1<br>[1]   |   |  |
| <b>(ii)</b> $\frac{dy}{dx} = \frac{3}{x} + 1 - 2x$<br>$\Rightarrow$ At R, $\frac{dy}{dx} = 0 = \frac{3}{x} + 1 - 2x$<br>$\Rightarrow 3 + x - 2x^2 = 0$<br>$\Rightarrow (3 - 2x)(1 + x) = 0$<br>$\Rightarrow x = 1.5$ , (or $-1$ )<br>$\Rightarrow y = 3 \ln 1.5 + 1.5 - 1.5^2 = 0.466$ (3 s.f.)<br>$\frac{d^2y}{dx^2} = -\frac{3}{x^2} - 2$<br>When $x = 1.5$ , $d^2y/dx^2 (= -10/3) < 0 \Rightarrow \max$ | M1<br>A1cao<br><br>M1<br>M1<br>A1<br>M1<br>A1cao<br><br>B1ft<br><br>E1<br>[9] | $d/dx (\ln x) = 1/x$<br><br>re-arranging into a quadratic = 0<br>factorising or formula or completing square<br>substituting their $x$<br><br>ft their $dy/dx$ on equivalent work<br><br>www – don't need to calculate 10/3   | SC1 for $x = 1.5$ unsupported, SC3 if verified<br><br><br><br>but condone rounding errors on 0.466 |
| <b>(iii)</b> Let $u = \ln x$ , $du/dx = 1/x$<br>$dv/dx = 1$ , $v = x$<br>$\Rightarrow \int \ln x dx = x \ln x - \int x \cdot \frac{1}{x} dx$<br>$= x \ln x - \int 1 dx$<br>$= x \ln x - x + c$<br><br>$\Rightarrow A = \int_1^{2.05} (3 \ln x + x - x^2) dx$<br>$= \left[ 3x \ln x - 3x + \frac{1}{2}x^2 - \frac{1}{3}x^3 \right]_1^{2.05}$<br>$= -2.5057 + 2.833..$<br>$= 0.33$ (2 s.f.)                  | M1<br><br>A1<br><br>A1<br><br>B1<br><br>B1ft<br><br>M1dep<br>A1cao<br>[7]     | parts<br><br><br>condone no $c$<br><br>correct integral and limits (soi)<br><br>$\left[ 3 \times \text{their } 'x \ln x - x' + \frac{1}{2}x^2 - \frac{1}{3}x^3 \right]$<br>substituting correct limits dep 1 <sup>st</sup> B1 | allow correct result to be quoted (SC3)  |

|  |   |   |  |
|--|---|---|--|
| <b>9(i)</b> $(0, \frac{1}{2})$   | B1<br>[1]   | allow $y = \frac{1}{2}$ , but not $(x =) \frac{1}{2}$ or $(\frac{1}{2}, 0)$<br>nor $P = 1/2$  |  |
| <b>(ii)</b> $\frac{dy}{dx} = \frac{(1+e^{2x})2e^{2x} - e^{2x} \cdot 2e^{2x}}{(1+e^{2x})^2}$<br>$= \frac{2e^{2x}}{(1+e^{2x})^2}$<br>When $x = 0$ , $dy/dx = 2e^0/(1+e^0)^2 = \frac{1}{2}$   | M1<br>A1<br>A1<br><br>B1ft<br>[4]                                 | Quotient or product rule<br>correct expression – condone missing bracket<br>cao – mark final answer<br><br>follow through their derivative  | product rule: $\frac{dy}{dx} = e^{2x} \cdot 2e^{2x}(-1)(1+e^{2x})^{-2} + 2e^{2x}(1+e^{2x})^{-1}$<br>$-\frac{2e^{2x}}{(1+e^{2x})^2}$ from $(udv - vdu)/v^2$ SC1 |
| <b>(iii)</b> $A = \int_0^1 \frac{e^{2x}}{1+e^{2x}} dx$<br>$= \left[ \frac{1}{2} \ln(1+e^{2x}) \right]_0^1$<br><i>or</i> let $u = 1 + e^{2x}$ , $du/dx = 2e^{2x}$<br>$\Rightarrow A = \int_2^{1+e^2} \frac{1/2}{u} du = \left[ \frac{1}{2} \ln u \right]_2^{1+e^2}$<br>$= \frac{1}{2} \ln(1+e^2) - \frac{1}{2} \ln 2$<br>$= \frac{1}{2} \ln \left[ \frac{1+e^2}{2} \right]^*$   | B1<br><br>M1<br>A1<br><br>M1<br><br>A1<br><br>M1<br><br>E1<br>[5] | correct integral and limits (soi)<br><br>$k \ln(1 + e^{2x})$<br>$k = \frac{1}{2}$<br><br><i>or</i> $v = e^{2x}$ , $dv/dx = 2e^{2x}$ o.e.<br><br>[ $\frac{1}{2} \ln u$ ] or [ $\frac{1}{2} \ln(v + 1)$ ]<br><br>substituting correct limits<br><br>www | condone no dx<br><br><br><br><br><br><br><br><br>allow missing dx's or incompatible limits, but penalise missing brackets                                      |
| <b>(iv)</b> $g(-x) = \frac{1}{2} \left[ \frac{e^{-x} - e^x}{e^{-x} + e^x} \right] = -\frac{1}{2} \left[ \frac{e^x - e^{-x}}{e^x + e^{-x}} \right] = -g(x)$<br>Rotational symmetry of order 2 about O   | M1<br>E1<br><br>B1<br>[3]   | substituting $-x$ for $x$ in $g(x)$<br>completion www – taking out $-ve$ must be clear<br>must have ‘rotational’ ‘about O’, ‘order 2’ (oe)  | not $g(-x) \neq g(x)$ . Condone use of f for g.  |
| <b>(v)(A)</b> $g(x) + \frac{1}{2} = \frac{1}{2} \cdot \frac{e^x - e^{-x}}{e^x + e^{-x}} + \frac{1}{2} = \frac{1}{2} \cdot \left( \frac{e^x - e^{-x} + e^x + e^{-x}}{e^x + e^{-x}} \right)$<br>$= \frac{1}{2} \cdot \left( \frac{2e^x}{e^x + e^{-x}} \right)$<br>$= \frac{e^x \cdot e^x}{e^x(e^x + e^{-x})} = \frac{e^{2x}}{e^{2x} + 1} = f(x)$<br><b>(B)</b> Translation $\begin{pmatrix} 0 \\ 1/2 \end{pmatrix}$<br><b>(C)</b> Rotational symmetry [of order 2] about P | M1<br>A1<br><br>E1<br>M1<br>A1<br><br>B1<br>[6]                   | combining fractions (correctly)<br><br><br><br>translation in y direction<br>up $\frac{1}{2}$ unit dep ‘translation’ used<br>o.e. condone omission of $180^\circ$ /order 2  | allow ‘shift’, ‘move’ in correct direction for M1.<br>$\begin{pmatrix} 0 \\ 1/2 \end{pmatrix}$ alone is SC1.   |